PUDOVIK, A. N.; MOSHKINA, T. M.; KHRAMTSOVA, V. P.

Diazophosphinic and hydrazediphosphinic esters. Zhur. ob.
khim. 33 no.1:94-97 '63.   (MIRA 16:1)

1. Kazanskiy filial Nauchno-issledovatel'skogo kinofotoinstituta.

(Phosphinic acid)   (Diazo compounds)
(Hydrazo compounds)

KHRAMUSHIN, A.

DANILENKO, A.; CHUMAKOV, N.:SERBINOVSKIY, G.; GRACHEV, V.; KHRAMUSHIN, A.;
        SOKOLOV, B.; BOL'SHAM, Ya.;TAYTS, A.; NEYFEL'D, M; FRENKEI', S.;
LYUDMIRSKIY, I.; NEBESNYY, A.; VESHENEVSKIY, S.; YERMILOV, A.;
BROZGOL', M.; SOLOV'YEV, P.; KLYUYEV, S.; ROZENTAL', A.;SMIRNOV,V.:
DOROFEYUK, A.

Solomon Mikhailovich Livshits; obituary. Prom energ. 11 no.12:34
D '56.   (MIRA 10:1)
            (Livshits, Solomon Mikhailovich, 1901-1956)

BELOV, N.N.; BOL'SHAM, Ya.M•; GORDEYEV, A.N.; GRACHEV, V.A.; YERMILOV, A.A.;
ZALESSKIY, A.M.; KIZEVETTER, Ye.N.; KNORRING, G.M.; KONSTANTINOV,
B.A.; KOPYTOV, N.V.; LEVIT, G.O.; MILLER, G.P.; NAYFEL'D, M.P.;
PRINTSEV, A.A.; SERBINOVSKIY, G.V.; SOKOLOV, B.A.; STASILOYTS, A.B.;
TAYTS, A.A.; KHRAMUSHIN, A.M.

Mikhail Konstantinovich Kharchev; obituary. Belov and others. Prom.
energ. 12 no.12:33 D '57.   (MIRA 10:12)
            (Kharchev, Mikhail Konstantinovich, 1896-1957)

YAKUBOVSKIY, F.B., red.; BELYAYEV, B.I., red.; VOLMYANSKIY, A.K., red.;
    KAMINSKIY, D.N., red.; KOL'TSOV, A.G., red.; KUREK, N.M., red.;
    OVSYANKIN, V.I., red.; PRIVALOV, N.N., red.; KHRAMUSHIN, A.M.,
    red.; ERISTOV, V.S., red.; UDOD, V.Ya., red.izd-va; TEMKINA,
    Ye.L., tekhn.red.

[Papers and reports of the section on industrial construction,
assembling and specialized work of the All-Union Conference on
Construction] Doklady i soobshcheniia. Moskva, Gos.izd-vo lit-ry
po stroit., arkhit. i stroit.materialam, 1958. 438 p.
                                                    (MIRA 12:7)

1. Vsesoyuznoye soveshchaniye po stroitel'stvu. Moscow, 1958.
Sektsiya promyshlennogo stroitel'stva, montazhnykh i spetsializi-
rovannykh rabot.
                        (Building)

KHRA muslin/l.)))

KHRAMUSHIN, A.M., inzh.; FRENKEL', S.N., inzh.

Growth of designing and installation organizations concerned with the electrification of industry in the U.S.S.R. Nov. tekh. i pered. op. v stroi., no.11:18-24 N '57.   (MIRA 10:12)
(Electrification)

1. YAKUBOV, M.K.: KHRAMUSHINA-PUSHKAR', L.M.

2. USSR (600)

4. Oils and Fats

7. Processing and refining of soap stocks. Masl. zhir. prom. 17. no. 9. 1952.

9. **Monthly** **List** **of** **Russian** **Accessions**, Library of Congress, February 1953. Unclassified.

KHRAMUSHINA-PUSHIKAR, L. M.

Improving the Quality of the Black Cotton Soap Stock with a Concentrated
Base. Leka Promishlenost (Light Industry), #7-12:28:July-Dec 1955

USSR.

Refining of black cottonseed soap stock with concentrated lye. M. K. Yakubov and L. M. Khramushina-Pushkar (Polytech. Inst., Kharkov).—*Maslobolno-Zhirovaya Prom.* 20, No. 3, 14-16(1955).—Black cottonseed soap stock contg. 30-40% fatty acids is mixed with 40-60% soln. of NaOH (20% in excess of the amt. required to saponify the neutral fat) and then heated to 170-250°. The authors claim that palmitic and acetic acids are formed from oleic, as in the Varentrap's reaction (KOH), and that the resulting soap is hard and light in color. Vladimir N. Krukovsky

BALAKIN; Ye.D.; MUKHANOV, G.V.; MURVANIDZE, D.S., red.; KHRAMYKH, N.M., red.; BRODSKIY, V.S., tekhn. red.

[Topics for inventors and efficiency promoters in the shoe industry]Temnik dlia izobretatelei i ratsionalizatorov obuvnoi promyshlennosti. Moskva, Biuro tekhn. informatsii legkoi promyshl., 1959. 95 p. (MIRA 15:11)

1. Russia (1923- U.S.S.R.)Gosudarstvennyy nauchno-tekhnicheskiy komitet.

(Shoe manufacture--Technological innovations)

KHAMRAEAYEV. I.Kh.; KHRAMYSHKIN, P.V.; RAKHMATULLAYEV, Kh.R.; RUZMATOV, S.R.

Geological and geochemical correlation of pyrite manifestations in the central Kyzyl Kum with some pyrite deposits in the Southern Urals. Uzb. geol. zhur. 8 no.4:19-29 '64.

(MIRA 18:5)

1. Institut geologii i geofiziki imeni Abdullayeva AN UzSSR.

KHRANIKOV,P.I., inzhener

Burning cut peat. Tekst.prom.8 no.2:40-42 F '48. (MIRA 8:11)
(Peat)

KHRANILOV, P. I.

36242

Progressivnyye normy raskhoda para (N_a sushku osnory). Tekstil. prom-st',
1949, No. 11, s. 40-41

SO: Letopis' Zhurnal'nykh Statey, No. 49, 1949

KHRANILOV, P.I.

Evaluation of the economic aspects of drying apparatus.
Izv. vys. ucheb. zav.; tekh. tekst. prom. no.4:116-121
'63.   (MIRA 16:11)

1. Ivanovskiy tekstil'nyy institut imeni M.V. Frunze.

KHRANILOV, P.I.

KHRANILOV, P.I., inzhener.

Increasing the productivity of drying apparatus. Tekst.prom. 14
no.5:36-39 My '54.   (MIRA 7:6)
        (Drying apparatus)

KHRANILOV, P.I.

Secondary power sources of finishing plants. Tekst.prom. 14 no.9:
36-37 S '54.   (MLRA 7:11)
    (Textile finishing)

KHRANILOV, P.I.; RAGOZINA, N.M.; SMIRNOVA, F.K.

Drying of fabrics under longitudinal blowing by the drying agent
at high velocities. Izv.vys.ucheb.zav.; tekh.tekst.prom. no.1:
134-138 '62.   (MIRA 15:3)

1. Ivanovskiy tekstil'nyy institut im. M.V.Frunze.
                        (Textile fabrics--Drying)

KHRANILOV, P.I.

Drying of fabrics by means of cold air. Izv.vys.ucheb.zav.;
tekh.tekst.prom. no.2:139-142 '63.   (MIRA 16:6)

1. Ivanovskiy tekstil'nyy institut imeni M.V.Frunze.
                (Textile fabrics—Drying)

KHANICOV, I.I.; DAGOLIN, N.M.

Intensification of the drying of fabrics by the speed increase
of the blower. Izv. vys. ucheb. zav.; tekh. teks. prom. no.3:
123-129 '64.   (MIRA 17:10)

1. Ivanovskiy tekstil'nyy institut imeni Frunze.

KHRANILOV, P.I., dotsent

Cases when gas dryers are more economical than steam dryers.
Tekst.prom. 25 no.1:83-84 Ja '65.   (MIRA 18:4)

1. Ivanovskiy tekstil'nyy institut.

L 06341-67   EWT(m)/EWP(t)/ETI   IJP(c)   JH/MJW/JD/JG/WB

ACC NR: AP6030322   (A)   SOURCE CODE: UR/0153/66/009/003/0449/0452

AUTHOR: Khranilov, Yu. P.; Poroykova, V. S.

ORG: Department of Technology of Electrochemical Manufactures, Ivanovo Chemical Engineering Institute (Kafedra tekhnologii elektrokhimicheskikh proizvodstv, Ivanovskiy khimiko-tekhnologicheskiy institut)

TITLE: Corrosion and anodic behavior of Mg-Al-Li alloys in concentrated sulfuric acid

SOURCE: IVUZ. Khimiya i khimicheskaya tekhnologiya, v. 9, no. 3, 1966, 449-452

TOPIC TAGS: corrosion, magnesium alloy, aluminum alloy, lithium alloy, SULFURIC ACID, ELECTRIC POTENTIAL

ABSTRACT: The purpose of the study was to determine whether the introduction of certain amounts of aluminum into a magnesium-lithium alloy would make its potential more electronegative in concentrated sulfuric acid (20-26 N H_2SO_4). The alloys were prepared from Mg-1 magnesium, AV-000 aluminum and LE-1 lithium, cast in a cylindrical mold, and subjected to homogenizing annealing in nitrogen for 40 hr at 300°C. The introduction of 1-5% Al into the Mg-Li alloy was found to raise the negative potential sharply, as expected. This alloy can therefore be used as the anodic material in activated chemical sources of current with lead dioxide cathodes. Most interesting are Mg-base alloys containing 5-6% aluminum and 10-17% lithium, which have a high negative

Card 1/2   UDC: 669.721.5'71'884:620.193.41

L 06341-67
ACC NR: AP6030322

potential and a low corrosion rate. Orig. art. has: 1 figure and 2 tables.

SUB CODE: 11,07/ SUBM DATE: 18Jul64/ ORIG REF: 004

6

Card   2/2

L 21127-65   EWT(m)/EPF(n(-2/EWA(d)/EWP(t)/EPR/EWP(b)   Ps-4/Pu-4   IJP(c)   JD/
JG/WB
ACCESSION NR: AP5001755   S/0153/64/007/005/0810/0815

AUTHOR: Poroykova, V. S.; Khranilov, Yu. P.; Sokolova, A. G.

TITLE: Corrosion resistance and anodic behavior of Mg-Li alloys in
concentrated sulfuric-acid solutions

SOURCE: IVUZ. Khimiya i khimicheskaya tekhnologiya, v. 7, no. 5,
1964, 810-815

TOPIC TAGS: magnesium alloy, lithium containing alloy, alloy micro-
hardness, alloy corrosion resistance, alloy polarization, alloy
potential

ABSTRACT: Magnesium-lithium alloys containing up to 20% lithium
have been studied. The microhardness of the alloy increased with in-
creasing lithium content in α-solid solution but dropped with the
appearance of the β-phase. Resistivity continously increased with
increasing lithium content, first sharply and then more slowly. The
atmospheric corrosion resistance of alloys containing up to 1.26% lithium was
found to be somewhat higher than that of unalloyed magnesium, but at

Card 1/3

L 21127-65
ACCESSION NR: AP5001755

higher lithium content corrosion resistance drops. All the alloys
corroded heavily in H_2SO_4 solutions, but more so in solutions with a
medium concentration than in 18—26-N solutions. In 18-N sulfuric acid,
α-alloys have the highest electrode potential, close to that of magne-
sium, but they polarize less than magnesium. Alloys of the α+β-
region have 0.1—0.15 v more negative potential than that of α-alloys
and are polarized at higher current density. The β-alloys have an
even more negative potential, ranging from -1.4 to -1.55 v, but are
easily polarized. After a certain period, potentials of all alloys
attain a steady value which is maintained for a considerable time (up
to 60 min). The potential height depends upon the kinetics of forma-
tion of oxide film on the surface of alloys, which in turn depends on
the number of cathodic and anodic areas on the surface. It is noted
that all the alloys including magnesium have a positive differential
effect in concentrated sulfuric acid. Orig. art. has: 7 figures.

ASSOCIATION: Ivanovskiy khimiko-tekhnologicheskiy institut (Ivanov
Chemical Technological Institute)

Card 2/3

L 21127-65
ACCESSION NR: AP5001755   0

SUBMITTED: 31Aug63   ENCL: 00   SUB CODE: MM, GC

NO REF SOV: 004   OTHER: 001   ATD PRESS: 3165

Card 3/3

TURNOV, B.S.; VINOGRADOV, P.A.; DOLGOPLOSK, B.A.; KHRANINA, Ye.N.; KOSTINA, S.I.

Effect of ethers on the chain structure in the stereospecific poly-
merization of butadiene. Dokl. AN SSSR 146 no.5:1141-1142 O '62.

(MIRA 15:10)

1. Yaroslavskiy zavod sinteticheskogo kauchuka. 2. Chlen-korrespon-
dent AN SSSR (for Dolgoplosk).
(Ethers)   (Butadiene)   (Polymerization)

KHRANOV,K., k.t.n. inzh.; DRAGUNCHEV, K., khimik; PASHOV,P.

Purification of natural water from manganese under semiopera-
tional conditions. Khidrotekh i melior 8 no.3:90-92 '63.

KICHEV Kiril, a tlo archus TUCHEV, Tsacho, inzn.

Struma River, and pollution of its water. Khidrotekh i melior
9 no.8-9:1-234 '64.

KHRANOV, L.S., professor.

Automatic leveling instrument. Nauka i zhizn' 24 no.3:
55 Mr '57.   (MLRA 10:5)
(Leveling)

KHRANOV, L.S., prof.; PRESNYAKOV, A.

Midget calculating machines. Nauka i zhizn' 26 no.2:73 F '59.
(MIRA 12:2)
(Calculating machines)

KHRANOVA, A.

Etiology of endemic goiter. Izv. med. inst., Sofia 8:115-144 1953.

(CLML 25:5)

1. Doctor.

KHRANOVA, A.

Studies on the effect of carbon dioxide in water and of humidity on development of experimental goiter in rats. Izv. med. inst., Sofia 8:159-170 1953.   (CIML 25:5)

1. Doctor. 2. Study made with the collaboration of Prof. Dr. Mosko Moskov, for histology and Mariya Andreycheva Vankova for chemistry.

KHRANOVA, A.

Influence of food with a large quantity of cooking salt (NaCl) and
sugar on the conditioned reflex action in rats. Izv biol med. BAN
3 no.2:31-39 '59.   (EEAI 10:4)

1. Institut po eksperimentalna meditsina pri Bulgarskata akademiia
na naukite.
        (FOOD)
        (SALT)
        (SUGAR)
        (CONDITIONED RESPONSE)

KHRANOVA, A.

(4)

Sofia, Doklady Bolgarskay Akademii Nauk, Vol 14, No 4, 1951 (continued)

24. "The Influence of [Experiment] on the Development of Experimental Goitre in Man," (A. Khranova, P. [...], and I. [...] (in Russian with French Summary); pp [...].

25. "The Influence of a Salt-Rich and Salt-Poor Diet on Blood Pressure in Rats of Various Ages," A. Khranova (in Russian with German Summary); pp 635-640.

KHRANOVA, A.; MOSKOV, M.; VASILEV, I.

The white cheese as preventive means against the development of
goiter in rats. Doklady BAN 14 no.6:655-658   '61.

1. Institut fiziologii Bolgarskoy Akademii nauk. Predstavlene akad.
D. Orakhovats.

KHRANOVA, A.

The influence of table salt and sugar in the diet on the blood pressure of rats of different ages. Doklady BAN 14 no.4:425-28 '61.

1. Institut fiziologii Bolgarskoy Akademii nauk. Predstavleno akad. D. Orakhovats.

KHRANOVA, A.; MOSKOV, M.; VASILEV, I.

Influence of the food for the development of experimental
goiter in rats. Doklady BAN 14 no.4:421-424 '61.

1. Predstavleno akad. D. Orakhovats.

KHRANOVA, A.

Effect of diets rich in kitchen salt (NaCl) and alternatively in kitchen salt and sugar on blood pressure in various age groups. II. Izv. inst. fiziol. 5:119-132 '62.

(SODIUM CHLORIDE nutrition & diets)
(CARBOHYDRATES nutrition & diets)
(BLOOD PRESSURE)   (AGING)

KHRANOVA, A.; TSANKOVA, E.

Importance of vitamins B1 and C for preservation of normal blood
pressure following higher nervous activity excitation in rats.
Dokl. bolg. akad. nauk 15 no.6:661-664 '62.

1. Predstavleno akad. D. ~~Kanate~~
            (THIAMINE)   (ASCORBIC ACID)   (CENTRAL NERVOUS SYSTEM)
                (BLOOD PRESSURE)   (STRESS)

KHRANOVA, A.; TSANKOVA, E.

Effect of overstraining of higher nervous activity on blood
pressure in the rat with dietary addition of kitchen salt
(NaCl) and resp. vitamins. Izv. inst. fiziol. (Sofiia) 6:
115-122 '63.

(BLOOD PRESSURE)   (ASCORBIC ACID)
(THIAMINE)   (PHARMACOLOGY)   (SODIUM CHLORIDE)
(CENTRAL NERVOUS SYSTEM)   (PHYSIOLOGY)
                    (STRESS)

ORAKHOVATS, D. [deceased]; KHRANOVA, A.; TSANKOVA, E.

The importance of diets with special reference to the content of proteins (animal and vegetable) and vitamins in circulatory dynamics (blood pressure) in rats exposed to overstraining of higher nervous activity. Izv. Inst. fiziol. (Sofiia) 7:81-94 '64.

OVAKHOVATS, D. [deceased]; KIRANOVA, A.

The importance of diets with special reference to the content of proteins (animal and vegetable) and vitamins in circulatory dynamics (blood pressure) in rats exposed to overstraining of higher nervous activity. II. Izv. Inst. fiziol. (Sofiia) 7: 95-115 '64.

KHRANOVA, A.

Effect of foods rich in eggs boiled during various periods on the blood pressure in rats with overstrained higher nervous activity and fed diets with and without vitamin C. Izv. inst. fiziol. (Sofiia) 8:49-58 '64

Effect of vitamin C on the blood pressure in protein-deficient rats with overstrained higher nervous activity. Ibid.8:9-67

KHRANOVA, V. N.

V. N. Khramova and A. P. Kreshkov, "Raising the Cement Frost-resisting Properties."

Report presented at the Second All-Union Conference on the Chemistry and Practical Application of Silicon-Organic Compounds held in Leningrad from 25-27 September 195?.
Zhurnal prikladnoy khimii, 1959, Nr 1, pp 238-240 (USSR)

ALEKSEYEV, A.P., otv. red.; ADROV, M.M., spets. red.; KONSTANTINOV,
        K.G.. spets. red.; KUTAKOV, B.G., red.; MASLOV, N.A., red.;
        MINDER, L.P., red.; NIKOL'SKIY, L.S., red.; STAROVOYTOV,
        P.A., red.; SURKOV, S.S., red.; KHRANOVSKIY, A.Yu., red.;
        YUDANOV, I.G., red.; VOROB'YEV, A.T., red.

        [Materials of the session of the Scientific Council of the
        Arctic Scientific Research Institute of Marine Fisheries
        and Oceanography dealing with the results of research in
        1962-1963] Materialy sessii Uchenogo soveta PINRO po rezul'-
        tatam issledovanii v 1962-1963 gg. Murmansk, 1964. 237 p.
                                            (MIRA 18:1)
        1. Murmansk. Polyarnyy nauchno-issledovatel'skiy i proyekt-
        nyy institut morskogo rybnogo khozyaystva i okeanografii.
        2. Direktor Polyarnogo nauchno-issledovatel'skogo i proyekt-
        nogo instituta morskogo rybnogo khozyaystva i okeanografii,
        Murmansk (for Alekseyev). 3. Laboratoriya vosproizvodstva
        Polyarnogo Nauchno-issledovatel'skogo i proyektnogo instituta
        morskogo rybnogo khozyaystva i okeanografii, Murmansk (for Sur-
        kov). 4. Laboratoriya tekhniki promyshlennogo rybolovstva
        Polyarnogo nauchno-issledovatel'skogo i proyektnogo instituta
        morskogo rybnogo khozyaystva i okeanografii, Murmansk (for
        Starovoytov).

KHRANOVSKIY, P.A.

KHRANOVSKIY, P.A.

4696 Kravchenko, N.A., Kravets, GKI, Khranovskiy, P.A. Iskusstvennyy
otbor zhivotnykh. Posobiye dlya uchiteley. Kiyev, u rad. Shkola", 1954.
160 s. s ill. 23 sm. 5000 ekz. 3 r 90k V per.- NA Ukr Yaz.- (54-58283) 636.082

SO: Letopis' Zhurnal nyph Statey, Vol 7, 1949

KHRANOVS'KIY, P.A.

      Practice in developing methods for the homogeneous transplantation
of sexual glands in domesticated water fowl.   Nauk.zap.Kiev.un. 13 no.
6:111-122 '54.   (MLRA 9:10)

      (Generative organs--Transplantation)
      (Water birds)

KHRANOVSKIY, P.A.

Importance of the size variability of the sexual glands in selecting breeds of domestic fowl. Dokl.AN SSSR 104 no.3:482-484 S '55.
(MLRA 9:2)

1.Kiyevskiy gosudarstvennyy universitet imeni T.G.Shevchenko.
Predstavleno akademikom Ye. N., Pavlovskim.
(Poultry breeding)

USSR/Farm Animals. Domestic Birds   Q-5

Abs Jour : Ref Zhur - Biol., No 11, 1958, No 50105

Author   : Khranovskiy, P.A.
Inst   : University of Kiev
Title   : Some Data on Size Variability of Sex Glands in Geese

Orig Pub : Nauk. zap. Kiivs'k. un-t, 1956, 15, No 11, 75-81

Abstract : In 1953, the author collected some data on the correlation
            between the size of sex glands in local breed geese and
            fertility. Subsequently, 700 birds were examined, whereby
            surgical methods (laparotomy) were used. The length and
            width of testes and ovaries in geese of various breeds
            (Barkov, Ramen, and Chinese) were measured. Size variabil-
            ities found in sex organs of geese whose breeds were com-
            pared here, were diverse. It was established that the
            variability in the size of testes is less pronounced. It
            was also found that testes in geese of Chinese breeds are
            larger, a fact which explains the higher fertility and
            greater sex activity of ganders of these breeds. The obtained

Card   : 1/2

USSR/Farm Animals. Domestic Birds   Q-5

Abs Jour : Ref Zhur - Biol., No 11, 1958, No 50105

            data prove that a direct correlation exists between the size
            of sex organs and their functions. -- A.P. Chmutova

Card   : 2/2

73

KHRANOVSKIY, P.A. [Khranovs'kyi, P.A.]

Morphological characteristics of the structure of the tail and
new theories of the evolution of exual asymmetry in birds.
Visnyk Kyiv.un. no.1. Ser.biol. no.2:113-121 '58.

(MIRA 16:4)

(SYMMETRY (BIOLOGY))   (TAIL)   (BIRDS—ANATOMY)

AUTHOR: Khranovskiy, P.A.   SOV/20-121-6-38/45

TITLE: On the Inheritance of Size and Function of the Reproductive
               Glands in Fowl (O nasledovanii razmerov i funktsii polovykh
               zhelez u domashnikh ptits)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol 121, Nr 6, pp 1083 - 1085
               (USSR)

ABSTRACT: At the moment scientists deal in genetics and selection of
               animals with the investigation of the regularities which de-
               termine variability and inheritance of the physical features.
               Almost all vital functions which determine the fundamental pro-
               ductive properties of animals are, however, connected with the
               inner organs and systems. In selection the investigations of the
               latter are, it is true, less useful. Nevertheless, this problem
               was already at issue at the beginning of our century (Refs 1-3).
               Also the first investigations of the inner organs were carried
               out (Refs 8-11). Publications reveal that inspite of the rapid
               development of research technique the selection methods for the
               breeding of domestic animals have to be completed. Since 1953
               the author has investigated the domestic goose with respect to
Card 1/3   the problem mentioned in the title; as we know this type of

On the Inheritance of Size and Function of the
Reproductive Glands in Fowl

animals still has a low index of fertility and of sexual pre-
cocity. In 4 generations (until 1957) 4000 geese have been in-
vestigated. Every year young ganders had to undergo laparotomy;
they were selected according to the degree of development of
their sexual glands. The selection amounted from 5-10%. Thus the
scientists succeeded in lengthening the sexual gland for 14,2 -
30 %. A detailed analysis showed that as a result of selection
in a great part of the ganders of the F_3 generation considerable

shifts in the transition of the glands to the active physiologi-
cal state set in: In 1953 among more than 500 ganders aged 3,5 -
4 months there was not a single animal with physiologically ac-
tive glands whereas among the F_3 birds (1956) there were up to

15 %. During the highest sexual activity the ganders had 30 %
larger testicles than the control birds. The physiological
changes with respect to fertility and sexual precocity correspond-
ed to this: In 1954 there were 21 % more fertilized eggs than in
the groups of negative selection. The sexual desire was observed
already earlier. The female offspring in further generations laid
eggs for the second time in summer. The total amount of eggs was

Card 2/3

On the Inheritance of Size and Function of the   SOV/20-121-6-38/45
Reproductive Glands in Fowl

              39 - 71 % higher. There are 11 references, 10 of which are Soviet.

ASSOCIATION: Kiyevskiy gosudarstvennyy universitet im. T.G. Shevchenko (Kiyev
               State University imeni T .G. Shevchenko)

PRESENTED: April 25, 1958, by Ye.N. Pavlovskiy, Member, Academy of Sciences,
               USSR

SUBMITTED: September 18, 1957

Card 3/3

KHRANOVSKIY, P.A.

Morphological features of the tail structure and new hypotheses
of the evolution of sexual asymmetry in birds. Zool.zhur. 38
no.12:1897-1899 D '59.   (MIRA 13:5)

1. Kiev State University.
    (Tail)   (Generative organs)   (Birds--Anatomy)

KHRANOVSKIY, P.A. [Khranovs'kyi, P.A.]

Sex, age and seasonal variations in the amount of erythrocytes
of domestic birds. Visnyk Kyiv. un. no. 5. Ser. biol. No.1:
88-97 '62.   (MIRA 16:5)

(POULTRY—PHYSIOLOGY)   (ERYTHROCYTES)

KHRAPACH, D. B.

Individual differences in the demand for vitamin B_1. I. The magnitude of the individual differences in the demand for vitamin B_1 among white rats. E. M. Maslenikova and D. B. Khrapach (Nutrition Inst., Acad. Med. Sci. U.S.S.R., Moscow). *Voprosy Pitaniya* **15**, No. 5, 8-19 (1956).— White rats were fed a vitamin B_1 (I)-deficient diet followed by the detn. of the excreted vitamin C (II) in the urine. Deficiency of I caused gradual disappearance of II in the urine; when the excretion of II ceased entirely I was added to the diet in the amt. necessary to restore the excretion of II in each of the 17 exptl. rats. The results indicate that the demand for I among individual rats to restore the excretion of II fluctuates between 5 and 75 γ/animal. Increase in the body wt. was usually observed with the restoration of the excretion of II or a few days earlier. The demand for I depends on metabolic activities and some special features of the higher nerve system among single animals.   E. W.

laboratorii izucheniya vitaminov

GRINBERG, Ye.A., dotsent; BABIY, Z.N.; LADUBA, T.L.; KHRAPACH, D.B.

Procurement of preserved blood in accomodations without special
equipment. Vrach. delo no.4:72-77 Ap'63.   (MIRA 16:7)

1. Kiyevskiy nauchno-issledovatel'skiy institut perelivaniya
krovi i neotlozhnoy khirurgii (nauchnyy rukovoditel' instituta-
prof. A.G.Karavanov).
          (BLOOD—COLLECTION AND PRESERVATION)

KHRAPACH, G.K.

Conversion of a turbocharged four-stroke engine from liquid to mixed fuel. Energ. biul. no.11:14-17 N '56.   (MLRA 9:12)
(Gas and oil engines) (Fuel)

KHRAPACH, Grigoriy Kuz'mich; MALYUKOV, G.A., red.; RASTOVA, G.V.,
   ved. red.

   [Installation and repair of compressors] Montazh i remont
   kompressorov. Moskva, Izd-vo "Nedra," 1964. 479 p.
                                           (MIRA 17:5)

KHRAPACH, G.K., inzh. (Kiyev)

The UKP20-60 transportable compressor installation for testing pipelines. Stroi. pred. neft. prom. 3 no.2:25-26 F '58.
(MIRA 11:4)

(Pipelines--Testing)
(Compressors)

KHRAPACH, Grigoriy Kuz'mich; MARTYNOVA, M.P., vedushchiy red.;
POLOSINA, A.S., tekhn.red.

[Compression plant operator; a manual] Mashinist kompressornoi
stantsii; spravochnoe rukovodstvo. Moskva, Gos.nauchno-tekhn.
izd-vo neft. i gorno-toplivnoi lit-ry, 1959. 307 p. (MIRA 12:9)
                    (Compressors)

ZAPARA, S.A., kand.tekhn.nauk; PANOV, V.A., inzh.; KHRAPACH, V.G., inzh.

Truck transportation in building Krivoy Rog Basin pits. Shakht.
stroi. 7 no.7:11-13 Jl '63.   (MIRA 16:10)

1. Nauchno-issledovatel'skiy gornorudnyy institut.

ZAPARA, S.A.; METS, Yu.S.; KHRAPACH, V.G.

Short-delay blasting and ways of increasing its efficiency in a
pit of the Krivoy Rog Southern Mining and Ore-dressing Combine.
Sbor. nauch. trud. NIGRI no.7:39-46 '60.   (MIRA 14:12)
            (Krivoy Rog Basin—Blasting)

KOMARISTOV, V.Ye., kand. tekhn. nauk, dots.; AVTUKHOV, I.V.,
        kand. tekhn. nauk, dots.; DUNAY, N.F., kand. sel'khoz.
        nauk, dots.; KHRAPACH, Ye.I., kand. tekhn. nauk;
        PESTRYAKOV, A.I., red.

        [Agricultural machines and implements] Sel'skokhoziai-
        stvennye mashiny i orudiia. [By] V.E.Komaristov i dr.
        Moskva, Kolos, 1964. 474 p.   (MIRA 17:12)

KHRAPACH, Ye.I., inzhener.

Effect of certain factors on the friction coefficient of straw
products. Sel'khozmashina no.8:5-7 Ag '57.   (MLRA 10:8)

1.Khar'kovskiy politekhnicheskiy institut imeni V.I. Lenina.
                        (Straw)

KHRAPACH, Ye.I., insh.

Calculating the strength of bale chambers in hay balers. Trakt. i
sel'khozmash. no.1:19-22 Ja '58.   (MIRA 11:4)

1.Khar'kovskiy politekhnicheskiy institut imeni V.I. Lenina.
                         (Hay--Harvesting)
                         (Agricultural machinery)

KHAYKIS, L., kand.tekhn.nauk; KHRAPACHEV, N., inzh.

Buckets with solid cutting lips. Stroitel' no.7:20 Jl '59.
(MIRA 12:10)
(Excavating machinery--Equipment and supplies)

KHAYKIS, L.B., kand.tekhn.nauk; KHRAPACHEV, N.N., inzh.

Repairing dippers with semicircular cutting edges. Transp.
stroi. 10 no.7:29-32 Jl '60.   (MIRA 13:7)
(Excavating machinery—Maintenance and repair)

KHRAPAL', A.A.

KHRAPAL', A.A. Sel'skoe khoziaistvo Aziatskogo Severa. Leningrad, Glavsevmorput',
1940.   131 p.   DA

SO: LC, Soviet Geography, Part I, 1951, Uncl.

KHRAPAL', A. A.

PA 33/49T4

USSR/Agronomy
Bibliography

Jul 48

"Eighth Grand Assembly of the Commission of the Far Northern All-Union Academy of Agricultural Sciences Imeni V. I. Lenin," A. A. Khrapal', Card Agr Sci, 2 pp

"Dok v-s Ak Selkhoz Nauk" No 7

Session convened in late Feb 48 was attended by representatives of far northern regions. All reports submitted were on achievements of agricultural science in regions where weather and climate are great obstacles. Delegates

USSR/Agronomy (Contd)

Jul 48

recommended measures which should be adopted by scientific research institutes to aid the polar farmer.

33/49T4

33/49T4

1. KHRAPAL',A. A.

2. USSR (600)

4. Geology and Geography

7. Transformation of Nature, Steppe and Forest Steppes, V. L. Kotel'nikov.
    (Moscow, Geography Press, 1949) Reviewed by A. A. Khrapal', Sov. Kniga,
    No. 12, 1949.

9. FDD Report U-3081, 16 Jan. 1953. Unclassified.

1. KHRAPAL', A. A.

2. USSR (600)

4. Kola Peninsula - Dairy Cattle

7. Organization of dairy cattle production on Kola Peninsula, Sots. zhiv., 15, no. 5, 1953.

9. Monthly List of Russian Accessions, Library of Congress, ___April___ 1953, Uncl.

Synthesis in the benzothiazole derivative series. A. I. Kiprianov and G. V. Klingel. *Uchenye Zapiski Khar'-kov. Univ.* 1940, No. 19, 205–11; *Khim. Referat. Zhur.* 4, No. 9, 68(1941).—A no. of derivs., including 2 monomethinecyanines, were obtained in expts. whose purpose was to study the acetylated derivs. of 2-mercapto-6-aminobenzothiazole as photographic stabilizers. 2-Mercaptobenzothiazole was used as the initial substance. A detailed scheme of the successive transformations and details of the synthesis are given. The following substances were obtained and described: 2-mercapto-6-aminobenzothiazole, m.: 264·5°; 2-mercapto-6-acetamidobenzothiazole, m. 208–5°; 2-methylmercapto-6-acetamidobenzothiazole, yellow crystals, m. 231–4°; 2-methylmercapto-6-benzamidobenzothiazole, m. 186–7°; (2-mercapto-6-benzothiazolyl)urea, m. 340° (decompn.); 1,3-bis(2-mercapto-6-benzothiazolyl)urea, m. 304–6°; 2-methylmercapto-6-aminobenzothiazole, m. 110–11°; (2-methylmercapto-6-benzothiazolyl)urea, m. 186–8°; 1,3-bis(2-methylmercapto-6-benzothiazolyl)urea, m. 238–40°; (2-methylmercapto-6-benzothiazolyl)thiourea, m. 163–4° (decompn.); (3-ethyl-6-acetamido-2-benzothiazole)(3-ethyl-2-benzothiazole)methinecyanine iodide, m. 205° (decompn.), max. of absorption 497 mμ; (3-ethyl-6-benzamido-2-benzothiazole)(3-ethyl-2-benzothiazole)methinecyanine iodide, m. 289° (decompn.), max. of absorption 498 mμ, crystallizes with 3 H$_2$O. A similar phenomenon is observed for thiocarbocyanines with substituting BzNH groups.

W. R. Henn

KHRAPAC, G.V.

Synthesis of some derivatives of β-phenylisopropylamine. I. Reaction of symmetric dihaloalkanes with β-phenyliso-propylamine. G. I. Kiprianov and B. E. Frenkel. *Ukrain. Khim. Zhur.* 16, No. 0, 620–6(1950)(in Russian).—Heating 5 g. $(CH_2Br)_2$ with 35 g. $PhCH_2CHMeNH_2$ (I) 5 hrs. at $130°$ gave 7.4 g. $(CH_2NHCHMeCH_2Ph)_2$, b_1 $105–200°$; di-HCl salt, m. $272–4°$. Heating 37.5 g. I with 17 g. $(CMe_2Br)_2$ 6 hrs. on a steam bath gave 5.6 g. $PhCH_2CHMeNHCMe_2CMe_2CH_2$, b_1 $135–43°$; HCl salt, m. $173–4°$; the free amine with Br yields a *product* m. $120–1°$. Heating 67.5 g. I with 12 g. 1,2-dibromocyclohexane 7 hrs. at $100°$ and 4 hrs. at $120°$ gave 7.2 g. N-(1-(or 2-)cyclohexen-1-yl)-2-phenyl-isopropylamine, b_1 $130–5°$; HCl salt, m. $180–9°$. $(CH_2)_3$-Br₂ (10.1 g.) and 67.5 g. I kept 24 hrs. at room temp. gave 10 g. 1,3-bis(2-phenylisopropylamino)propane, b_1 $200–5°$; di-HCl salt, m. $238–9°$. Similarly $(CH_2)_4Br_2$ gave 94% 1-(2-phenylisopropyl)pyrrolidine, b_1 $117–23°$; HCl salt, m. $130–8°$. $(CH_2)_5Br_2$ gave 63% 1-(2-phenylisopropyl)piperi-dine, b_1 $115–17°$; HCl salt, m. $215–16°$. $(CH_2)_6Br_2$ and I kept 24 hrs. at room temp. gave 58% 1,6-bis(3-phenyliso-propylamino)hexane, b_1 $224°$; picrate, m. $183–4°$; di-HCl salt, m. $253–4°$. II. Reaction of β-phenylisopropylamine with α-oxides. G. I. Kiprianov and G. V. Khrapal. *Ibid.* 627–31(1950)(in Russian).—Heating 10.5 g. ethylene oxide with 82 g. $PhCH_2CHMeNH_2$ (I) in a sealed tube 10 hrs. at $100°$ gave 25 g. N-(2-hydroxyethyl)-2-phenylisopropyl-amine, b_1 $117–21°$; sulfate, m. $200–5°$; HCl salt, m. $111–13°$. Propylene oxide similarly treated gave 66.5% N-(2-hydroxypropyl) homolog, b_1 $122–6°$, crystg. on standing and m. $65–4°$; HCl salt, m. $144–6°$. The residual *liquid portion* of the product gave a HCl salt, m. $142–5°$, which mixed with the previous HCl salt, m. $122–9°$, indicating different *stereoisomeric* forms. EtCH.CH₂O and I similarly gave

57.4% N-(2-hydroxybutyl) compd., b_1 $130–3°$; HCl salt, m. $115–30°$. Cyclohexene oxide and I after 70 hrs. at $110–60°$ gave 91% N-(2-hydroxycyclohexyl) analog, b_1 $148–53°$; HCl salt, m. $193–7°$. Glycidol and I after 2 hrs. at $160°$ gave 85% $PhCH_2CHMeNHCH_2CH(OH)CH_2OH$, b_1 $169–70°$; HCl salt, viscous mass. III. Reaction of β-phenylisopro-pylamine with 1,3-dibromopropanol. G. I. Kiprianov and G. V. Khrapal. *Ibid.* 632–8(1950)(in Russian).—HO-$CH(CH_2Br)_2$ (I) (8.7 g.) was added with ice cooling to 20 g. $PhCH_2CHMeNH_2$, the mixt. cooled 1 hr., kept 2 hrs. at room temp., acidified with HCl, extd. with Et_2O, and the acid soln. made alk. with NaOH and extd. with Et_2O, yielding 47.5% N-epoxyallyl-5-phenylisopropylamine (II), b_1 $143–5°$, and 20.2% 1,3-bis(2-phenylisopropylamino)-2-propanol (III), b_1 $215–20°$. II solidified on standing and m. $81–3°$; HCl salt, m. $107–8.5°$. III gave a *nitrate*, m. $170–1°$ (from H₂O), m. $166–7°$ (from EtOH); di-HCl salt, m. $247–50°$ (pure), m. $220–32°$ (crude). The yield of III can be raised by increasing the proportion of I in the syn-thesis. Keeping 4.11 g. epibromohydrin and 12.25 g. I 1 hr. gave 24.4% II and 47% III. Heating 2.6 g. II with 1.84 g. I 3 hrs. at $130–40°$ led to no reaction; at $175–85°$ 61% III was formed. II heated with 40% aq. $MeNH_2$ in sealed tube 6 hrs. at $170–80°$ gave 67.8% 1-methylamino-3-(2-iso-propylamino-2-propanol, b_1 $168–60°$; di-HCl salt, m. $235.5–6.5°$. II heated with excess EtNH₂ 5 hrs. at $170–80°$ gave 75.8% 1-diethylamino-3-(2-phenylisopropylamino)-2-pro-panol (IV), b_1 $155–60°$; HCl salt and sulfate, oils; oxalate, m. $145–7°$. N-Epoxyallyldiethylamine heated with excess I 9 hrs. at $175–85°$ gave 68% IV.   G. M. Kosolapoff

1.   KIPRYANO, G.I.;   KHRAPAL', G.V.

2.   USSR (600)

4.   Amines

7.   Synthesis of some derivatives of ss-phenylisopropylamine with a – oxicdes, G.I.
     Kipryanov, G.V. Khrapal', Ukr.khim.zhur. 16 no. 6, 1951.

9.   **Monthly List of Russian Accessions**, Library of Congress, ___APRIL___ 1953. Unclassified.

1. KIPRYANOV, G. I.; KHRAPAL', G. V.

2. USSR (600)

4. Oxides

7. Synthesis of some derivatives of β -phenylisopropylamine. Part 2. Reaction of β -phenylisopropylamine with α -oxides. Ukr. khim. zhur. 16, No. 6, 1951.

9. Monthly List of Russian Accessions, Library of Congress, ___April___ 1953, Uncl.

1. KIPRIANOV, G. I.; KHRAPAL', G. V.

2. USSR. (600)

4. Propanol

7. Synthesis of some derivatives of β-phenylisopropylamine. Part 3. Reaction of β-phenylisopropylamine with 1, 3-dibromopropanol, Ukr. khim. zhur., 16, no. 6, 1951.

9. Monthly List of Russian Accessions, Library of Congress, ___April___ 1953, Uncl.

KIPRIANOV, G.I.

KIPRIANOV, G.I.; KHRAPAL', G.V.

Synthesis of certain derivatives of β-phenylisopropyl-
amine. Part 5. N - β-chloroethyl and N - β-dimethyl-
aminoethyl derivatives. Uk.rkhim.zhur. 20 no.1:53-56 '54.
(MLRA 7:3)

1. Ukrainskiy institut eksperimental'noy endokrinologii, g. Khar'kov.
(Amines)

KHRAMOV, V.P.

18(6)

PHASE I BOOK EXPLOITATION SOV/3199

Akademiya nauk SSSR. Institut obshchey i neorganicheskoy khimii
im. N. S. Kurnakova

Analiz blagorodnykh metallov (Analysis of Noble Metals) Moscow,
1959. 193 p. Errata slip inserted. 2,700 copies printed.

Resp. Ed.: N. K. Pshenitsyn, USSR Academy of Sciences, Corre-
sponding Member; and O. Ye. Zvyagintsev, Doctor of Chemical
Sciences; Eds. of Publishing Houses: T. G. Levi, and D. N.
Trifonov; Tech. Ed.: I. N. Guseva.

PURPOSE: This collection of articles is for scientists engaged
in the study and analysis of the noble metals.

COVERAGE: This is a collection of articles on the analysis of the
noble metals. It includes studies carried out by the Institute
of General and Inorganic Chemistry im. N. S. Kurnakov (AN SSSR),
as well as reports presented by scientific research organizations
and by industrial enterprises at the Third and Fourth Conference
on Noble Metals held in 1954 and 1957, respectively. The
studies and reports describe new organic reagents for gravi-
metric determination of platinum metals, and physicochemical
methods of analysis (spectrophotometric, polarographic and
potentiometric). Special attention is given to spectral
analysis for the determination of admixtures in alloys of
platinum metals, silver, and gold, as well as in refined noble
metals. The collection also includes analytical methods, tables
and charts for material containing metals of the platinum
group, as well as a review of the literature on the analysis
of platinum metals published in the last five years. No
personalities are mentioned. References follow each chapter.

KHRAPAY, V. P. (NOVOSIBIRSK)

"Methods of Spectral Analysis in the Determination of Impurities in High-Purity Gold"

paper submitted to the Fifth Conference on the Analysis of Nobel Metals, Novosibirsk, 20-23 September 1960

So: Zhurnal analiticheskoy khimii, Vol XVI, No. 1, 1961, page 119

ABSTRACT: The author notes that after an arithmetic operation has been performed
on digital computers the result is normally rounded off. The magnitude of the re-
sultant error must be taken into account when determining the necessary number of
places for the representation of numbers in the machine. Generally speaking, esti-
mation of errors in addition and subtraction offers no particular difficulties;
the error in the product of two n-place numbers is also calculated with relative
ease in the event that the rounding-off operation is carried out after all 2n signi-
ficant digits of the product have been obtained. However, a number of methods of
multiplication can be performed in a more economical fashion if there is a prelim-
inary rounding-off of each product of the multiplicand and the multiplier digit.
In the present article, an estimate is given of the binary multiplication error
which arises in the event of this kind of rounding off-operation. The author pre-
Card 1/2

ACCESSION NR: AT4016490

sents 2 theorems and 7 lemma in a determination of the maximum and minimum values of δ. The problem of determining the required amount of additional places p for a given error in multiplication is also analyzed. Orig. art. has: 32 formulas and 1 table.

ASSOCIATION: none

SUBMITTED: 05Jul62   ENCL: 00

SUB CODE: DP   NO REF SOV: 001   OTHER: 000

n_b n δ

Card   2/2

AUTHOR: Khrapchenko, V. M.

TITLE: A method for the conversion of a multi-row into a single-row code

TEXT: A method is described for the fast conversion of an m-row code

$C = \sum_{j=1}^{m} A_j$ into a single-row code. Let the summands $A_j (1 \leqslant j \leqslant m)$ be

represented in a positional calculation system with the constant weight

$A_j = \sum_{i=1}^{n} a_{ji} r^{-i}$, where n is the number of the order of A_j. The integer r

is the basis of the system of counting applied $(r \geqslant 2)$, and a_{ji} is one of

the numbers $0, 1, \ldots, r-1$; $a'_{j'i'}$ equals $b_{j'(i'-(m'-j'))}$ for $1 \leqslant i'-(m'-j') \leqslant n$

and 0 for all other cases.

Card 1/3

A method for the conversion of a ...   S/020/63/148/002/014/037
B125/B112

$$C = \sum_{i'=1}^{m'} A'_{i'},$$

(8)

$$A'_{i'} = r^{m'-1} \sum_{i=1}^{n+m'-1} a'_{i'i''} r^{-i''},$$

is derived for the number C in the m'-row code. The number of rows in the code is reduced to two by repeated application of such steps. The two-row code is then converted into a single-row code by a parallel (n+m') digit summator. The optimum code conversion circuit of the computer can be chosen freely if the number m_k of the rows in the code (after the k-th step of conversion) is divided into p_k groups each of which has u_k summands, and into a residual group with v_k summands, according to $m_k = p_k u_k + v_k$ $(0 \leqslant v_k \leqslant u_k)$. The number of rows in these

Card 2/3

A method for the conversion of a ...   S/020/63/148/002/014/037
B125/B112

groups is reduced to two, and then to one by a summator. In this
variant no more than $(m-2)/(u-u')$ blocs are required, for a u'-row sum
of u single-row summands, no more than s less complex blocs and a one-bit
summator are required. Multiplying speed may be tripled or quadrupled by
the above method. There are 1 figure and 1 table.

ASSOCIATION: Institut elektronnykh upravlyayushchikh mashin Akademii
nauk SSSR (Institute of Electronic Controlling Engines of
the Academy of Sciences USSR)

PRESENTED: July 12, 1962, by A. I. Berg, Academician

SUBMITTED: July 12, 1962

Card 3/3

S/799/62/000/003/001/008

AUTHOR: Khrapchenko, V. M.

TITLE: On the synthesis of networks for the modeling of a disjunctive normal
             form.

SOURCE: Akademiya nauk SSSR. Institut elektronnykh upravlyayushchikh mashin.
             Tsifrovaya tekhnika i vychislitel'nyye ustroystva. no.3. 1962, 3-13.

TEXT: The problem of designing a minimal circuit of the "and-or" type to
realize certain algebraic-logic functions is reduced in this survey paper to its
simplification by means of a disjunctive normal form (DNF). A great number of
methods for the construction of a minimal DNF exists. The practically most signi-
ficant results obtained by a number of authors in this direction are selected. They
have been reduced to a unitary method for the design of a minimal DNF, which is
illustrated on the example of the design of a diode circuit. The example employed
in the discussion is an arbitrary two-stage diode circuit of the "and-or" type.
Here different variables and their inversions are placed into correspondence to the
various input signals of the circuit; it is assumed that an elevated signal level has
a corresponding variable or its inversion equal to "1" and a low level equal to "0".
Each group of diodes that form an "and" subcircuit will be juxtaposed to a conjunction
of variables and their inversions that correspond to the input signal of the diodes of
Card 1/2

On the synthesis of networks for the modeling S/799/62/000/003/001/008

that group; the entire DNF circuit, consisting of the conjunctions, will correspond
to its subcircuits. Obviously, between the "and-or" circuits and the DNF's, a
mutually single-valued correspondence is established, which exhibits the following
property: The level of the output signal of the circuit corresponds mutually single-
valuedly to the value of the function determined by the DNF in the set of the values
of the variables that correspond to the levels of the input signals. As an example,
a certain algebraic logic function, that depends on 3 variables, is assumed as given.
One of the DNF's of this function and its corresponding circuit are schematically
drawn. The reasonings derived from this example are developed into the above-
defined unitary design method. There are 1 figure, 6 tables, and 12 references
(5 Russian-language Soviet and 7 English-language, of which 1 in Russian transla-
tion).

Card 2/2

L   07925-67   EWT(m)/EWP(t)/ETI   IJP(c)   JD

ACC NR: AP6033385   SOURCE CODE: UR/0075/66/021/008/0980/0984

AUTHOR: Grushina, N. V.; Tsevun, V. I.; Khrapchenkova, G. V.;
Yerdenbayeva, M. I.; Kozin, L. F.

ORG: Institute of Chemical Sciences, AN KazSSR, Alma-Ata (Institut khimicheskikh
nauk AN KazSSR)

TITLE: Determination of impurities in high-purity cadmium

TOPIC TAGS: cadmium, cadmium metal, impurity determination, high purity
cadmium, cadmium nitrate

ABSTRACT: A method has been developed for the spectrochemical determination
of 10^{-4}—$10^{-6}\%$ impurities in cadmium after their concentration by coprecipitation
with cadmium diethyldithiocarbamate. The method was applied to the analysis
of high-purity cadmium metal and cadmium nitrate. The relative experimental
error is $\pm 25\%$. Orig. art. has: 2 figures and 3 tables. [Authors' abstract]

SUB CODE: 07/ SUBM DATE: 23Nov64/ ORIG REF: 007/ OTH REF: 001/

Card 1/1   vmb

KHRAPIN, T.M.

Oscillators instead of buzzers. Avtom., telem. i sviaz' 2 no.1:34 Ja '58.   (MIRA 11:1)

1. Starshiy elektromekhanik Grebenkovskoy distantsii Yuzhnoy dorogi.
(Oscillators, Electron-tube)   (Railroad--Telephone)

KHRAPIN, T.M.

Rectifier for intrastation communications. Avtom., telem. i sviaz'
2 no.6:34 Je '58.   (MIRA 11:6)

1. Starshiy elektromekhanik Grebenkovskoy distantsii signalizatsii
i svyazi Yushnoy dorogi.
            (Railroads--Communication systems--Equipment and supplies)

KHRAPKIN, M·A

Ca

PROCESSES AND PROPERTIES INDEX

Charge make-up of the eastern by-product coke plants.
N. S. Gryaznov and M. A. Khrapkin. *Stal* 7, 773-81
(1947).—An investigation was carried out for the purpose
of simplifying the compn. of coking charges and to adapt
coals hitherto not used for coking as substitutes for high-
grade coking coals. The coke from the new charges was
satisfactory for metallurgical use.　　　M. Hoech

21

ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION

KHRAPKIN, M. A.

FA 24T33

Sep. 1947

USSR/Engineering
    Coke Plants
    Fuels, Solid

"Construction of Mixing Sheds at Eastern By-product Coke Factories," N. S. Gryaznov, Candidate in Technical Sciences, M. A. Khrapkin, Engr, Eastern By-product Coke Institute, Main Coke Administration, 8½ pp

"Stal'" No 9

In accordance with the agreements of the All-Union Technical Conference on Coke in 1946, there has been much work on the reconstruction of eastern by-product coke factories and plants. This has decreased the number of components in the mixing sheds, led to the use of several new types of coal for making coke and increased the use of low-grade coke coal (particularly bituminous, gaseous, K Zh). The result was a standardization of mixing with a resultant standardization of the quality of coke. Well-illustrated with diagrams.

FA 24T33

KHRAPKIN, M.A., inzh.

Comparable technical and economic indices of the operations of coal preparation plants. Ugol' 36 no.11:42-44 N '61. (MIRA 14:11)
(Coal preparation plants)

A M KHRAPKO and V F RULEV

    "Development of a Device for Measuring the Degree of Coupling in
Magnetrons" from Annotations of Works Completed in 1955 at the State Union Sci.
Res. Inst; Min. of Radio Engineering Ind.

So: B-3,080,964

S/107/61/000/005/003/004
E192/E382

AUTHORS: Rasskazikhina, N. and Khrapko, R., Engineers

TITLE: An Underwater Communications Device ("Walkie-
            talkie")

PERIODICAL: Radio, 1961, No. 5, pp. 26 - 27

TEXT: The system described is an independent communications
device whose weight, together with supply sources, is about
1 kg. The equipment can be used for voice-communications
(talking) under water and consists of a throat microphone,
a low-frequency amplifier based on transistors, an underwater
loudspeaker and a supply source. Since a throat microphone is
used in this equipment, it can be employed by divers provided
with aqualungs or oxygen cylinders. The supply source is
provided by 8 or 10 pocket-lamp batteries (type КБС·Л·0.5
(KBS-L-0.5)). The useful life of a set of such batteries
is 30 - 50 min (continuous). The amplifier is shown in
Fig. 1, where the first stage is an emitter-follower which
provides a matching buffer between the microphone and the
input of the amplifier proper. The second stage is the first

Card 1/3

S/107/61/000/005/003/004

An Underwater Communications E192/E382

amplifier with a transformer load; this is followed by the
output stage, based on 2 transistors, type ∏4Б (P4V), which
operate in push-pull without an output transformer. The coil
of the loudspeaker, with a resistance of $5\,\Omega$, is the load to
the output amplifier. The output power of the amplifier is
25 W, its efficiency being about 75%. The underwater loud-
speaker is based on a dynamic loudspeaker, type 2ГД-3 (2GD-3)
(Fig. 2). The magnetic system 1 of the speaker is covered
by a duralumin ring 2 , which supports a diaphragm made of
celluloid, 0.3 - 0.5 mm thick. The diaphragm is fixed by
another ring 4 , which is screwed onto the top of the first
ring. The coil of the loudspeaker is fixed to a diaphragm and
its terminals are taken through small apertures in the
diaphragm which are hermetically sealed. The equipment is
switched off by means of the key K_1 , which is sealed inside
a small rubber bag.
There are 2 figures.

Card 2/3

KHRAPKO, R.I.

Uniqueness of a centrally symmetrical gravitational field
in a vacuum. Zhur.eksp. i teor.fiz. 49 no.6:1884-1886
D '65.   (MIRA 19:1)

1. Moskovskiy aviatsionnyy institut. Submitted July 5, 1965.

AUTHOR: Khrapko, R. I.

TITLE: Stationary quasi-onedimensional magneto-gasodynamic flow
              at finite conductivity

PERIODICAL: Zhurnal tekhnicheskoy fiziki, v. 32, no. 9, 1962, 1063-1071

TEXT: In order to study transitions from subsonic to supersonic flow
patterns a system of 9 equations is set up for the stationary quasi-
onedimensional movement of an ideal gas of constant finite conductivity σ
in a plane tube. The gas moves in x-direction, an electric field is
applied in y-direction, and an external magnetic field (not necessarily
homogeneous) is so directed that its z component is large compared with
its x and y components. The system of 9 equations contains the dependencies
of the following 11 functions on the x coordinate: Electric field strength
E_y, tube width A in y direction, electric current component j_y, gas
velocity v_x, density ϱ, pressure p, enthalpy w, velocity of sound a, total
magnetic field component H_z, external magnetic field component H_z^e,

Card 1/2

external field gradient $\partial H_x^e / \partial z$. Thus two additional conditions may be stipulated arbitrarily in order to treat special cases. The following cases are discussed: 1) A and H_z are constant. 2) A and the Mach number v/a are constant. 3) A and $H_z v/E_y$ are constant. 4) A is constant, and $\partial H_x^e / \partial z$ vanishes. 5) $\partial H_x^e / \partial z$ vanishes, and a is constant. It is shown how the system of equations can be generalized for the case where the conductivity also is variable. There are 3 figures. *M. F. Shirckev* *thanked for valuable discussions.*

SUBMITTED: June 12, 1961 (initially)
               December 18, 1961 (after revision)

Card 2/2

L 32628-66   EWT(1)   IJP(c)   WW/GG

SOURCE CODE: UR/0056/66/050/004/0971/0974

ACC NR: AP6014037

AUTHOR: Khrapko, R. I.

ORG: Moscow Aviation Institute (Moskovskiy aviatsionnyy institut)

TITLE: Propagation of light through a singular Schwartzschild sphere

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 50, no. 4, 1966, 971-974

TOPIC TAGS: cosmogony, light velocity, gravitation effect, general relativity theory, Doppler effect

ABSTRACT: The problem of propagation of light from an emitting surface of a gravitating sphere expanding with parabolic velocity from within its own singular sphere, in other words, how can an external observer see this surface, is solved in the present paper in a coordinate system with a metric

$$ds^2 = -\frac{r-1}{r}dt^2 + \frac{r}{r-1}dr^2 + r^2(d\theta^2 + \sin^2\theta\, d\varphi^2). \qquad (1)$$

To overcome the difficulty encountered in identifying parts of a geodesic which are separated by the singular sphere, the equation of the geodesic on one side of the singular sphere is regarded as the analytic continuation of this equation on the other side of the singular sphere. The equation of the world line of an emitter located on the surface of such a sphere is obtained by fixing the Lemaitre radio coordinate in the equation that gives the transition from the Schwarzschild system to the Lemaitre

Card 1/2

L 32628-66

ACC NR: AP6014037   2

system. Allowance is made for the Doppler shift of the frequency of light. Equations are derived which give the complete evolution of the picture seen by the observer, with the exception of the intensity. It is shown that the frequency ratio of the light at the edge of the disc and at the point of observation decreases monotonically from infinity (at the instant of the first flash). The frequency ratio of the light emitted by the luminous sphere as it passes through the singular sphere and received by a very distant observer is equal to 2. The light received by such an observer from the edge of the disc has its natural color at all times. The author thanks Professor M. F. Shirokov and V. Babetskiy for fruitful discussions. Orig. art. has: 1 figure and 8 formulas.

SUB CODE: 20, 03/   SUBM DATE: 11Oct65/   ORIG REF: 004/   OTH REF: 002

Card 2/2